


**Dr Z's Medical Coding Series:
Diagnostic & Interventional Cardiovascular Coding Reference: 2011
Edition**

2011 Book Errata



Text to be deleted has been crossed out and new text noted in blue font.

Page 292

CODING INSTRUCTIONS

15. CPT code 36002 is specific for thrombin injection to treat pseudoaneurysm. Use codes 37204/~~75984~~ **75894** when coil embolization is necessary. Use codes 34900/75954 when covered stent graft placement across an iliac pseudoaneurysm is required. Code 34900 is an inpatient only procedure (status indicator C) for Medicare.

Page 307

CODING INSTRUCTIONS

9. Coronary intervention rules dictate that if (regardless of reason) an angioplasty, atherectomy, and stent procedure are done in the same coronary vascular distribution, only the highest level of intervention (the stent procedure) can be coded. These ~~same~~ rules **differ for now apply to** infrainguinal arterial interventions, **as stent with atherectomy supersedes atherectomy, which supersedes stent placement, which supersedes angioplasty.**

10/6/2011

1

Page 310

EXAMPLES

- 1) Hypertensive patient undergoes abdominal aortogram with selective renal angiography of two left and one right renal arteries (36245-50, 36245-59, 75724-59, 75774). Severe stenoses of both left renal arteries are balloon dilated (~~35471-50~~, **35471, 35471-59**, 75966, 75968).
-

Page 326

CODING INSTRUCTIONS

11. There is a ~~hierachy~~ **hierarchy** with the lower extremity revascularization codes ~~similar to coronary interventions~~. Stent placement **with atherectomy** supersedes atherectomy, **which supersedes stent**, which supersedes angioplasty, when performed in the same vessel in the tibial/peroneal territory, or in any combination of vessels in the femoral/popliteal territory. Angioplasty is also bundled with stent placement in the same vessel in the iliac territory.

Page 326

CODING INSTRUCTIONS

18. The tibial/peroneal territory includes three vessels that are separately coded: the anterior tibial, posterior tibial, and peroneal arteries. The tibial/peroneal trunk is considered part of any distal vessel intervention **in the posterior tibial and peroneal arteries** (~~similar to the left main coronary artery~~). **The tibial/peroneal trunk is considered a separate vessel from the anterior tibial artery**. The dorsalis pedis is considered part of the anterior tibial artery, and the medial malleolar artery is considered part of the posterior tibial artery.

Page 328

CODING INSTRUCTIONS

33. If kissing stents (or angioplasty) are performed at the iliac bifurcation, one initial vessel and one additional vessel intervention as appropriate. If done in the femoral/popliteal territory (such as angioplasty in the profunda femoral and superficial femoral arteries extending back into the common femoral), only submit one femoral/popliteal code. **Note: The iliac bifurcation refers to the location where the common iliac artery divides into the internal and external iliac arteries.**

Page 329

CODING INSTRUCTIONS

51. Do code stent placement (37221, 37223, ~~37226, or 37227~~) for stent graft placement in an iliac ~~or femoral~~ vessel to treat stenosis (~~not aneurysm, pseudoaneurysm, AVM, or trauma~~). **Do code 37226 for stent graft placement in a popliteal artery to treat an aneurysm. In the iliac artery, use codes 34900 and 75954 for stent graft placement to treat an aneurysm.**

53. Ultrasound guidance for vascular access (76937) may be reported in addition to lower extremity revascularization procedures when there is medical necessity and all necessary elements are documented.

54. Do code catheter placement separately for TPA infusion in a lower extremity artery if the intervention (angioplasty/atherectomy/stent placement) is performed at a separate session on the same date or service or on a different date of service.

Page 358

CODES:

Second and third entries from the bottom should be

Iliac exposure with cutdown for graft delivery via conduit	◆33883 ◆34833	N/A	35.53			
Brachial exposure with cutdown to assist with graft delivery	◆34884 ◆34834	N/A	12.87			

Page 505

PROCEDURE

- The artery-narrowing drug ergonovine (~~or, alternatively, acetylcholine~~) is injected to provoke coronary artery spasm.