ERRATA for Interventional Radiology Coding Reference 2023 Edition

Text deletions are crossed out. Corrected text is **blue and bolded**. Ordered by appearance in text.

Page 204, Coding Instructions

5. Do not report PICC line codes for a <u>midline</u> catheter placement. Report code 36400, 36405, 36406, or 36410 as appropriate based on documentation.

Page 218, Code Table

Disruption of fibrin sheath with angioplasty balloon (via remote access)	36595 -52 -	5183	3.59	75901	N/A	0.49
	(or -74)					

Page 291, Coding Instructions

11. Do not code separately for selective catheter placement when using codes 37215, 37216, 37217, 37218, 0075T, 0076T, 34717, 34718, 61635, and 34841-34848, as these codes bundle the catheter placement related to the procedure. Non-selective catheter placements are bundled with codes 34701-34712, while selective catheter placements are reported with codes 34701-34712. Code separately for selective catheter placement when certain additional procedures are performed, such as stent placement in the renal artery (37236, 37237) or embolization of a non-target vessel (37242). Rules vary for other catheter placements depending on the type of endograft procedure performed (IBE vs. EVAR vs. FEVAR).

Page 447, Examples

1) Patient with a long-term **external** biliary catheter for benign strictures presents to the emergency room with his catheter in hand. In radiology, the site is prepped and draped and the exit site catheterized with a Christmas tree adapter through which contrast is injected, demonstrating a patent catheter tract with visualization of peripheral bile ducts (20501, 76080). Under fluoroscopic guidance using wires and catheters, access is gained into the biliary tree, and a new catheter is placed into the duodenum (add 47535; delete 20501, 76080).